



Low Socioeconomic Status is Associated with Adverse Events in Children and Teens on Insulin Pumps Under a Universal Access Program: A population-based Cohort Study

Rayzel Shulman MD, PhD


CAHSPR

May 10, 2016




+ Background and Rationale

- High incidence of type 1 diabetes (T1D) in Canada (25.9/100,000/year)
- The Ontario Pediatric Diabetes Network (PDN) oversees care delivery
- Differences in PDN center characteristics (2012):
 - 24-hour support
 - Centre type
 - Physician model of care
 - Nursing patient load
- Universal funding for pediatric insulin pump therapy in Ontario since 2006



PDN Current State Survey 2013

Pump Use Following the Introduction of a Universal Funding Program in 2006




+ Background and rationale

- Diabetic ketoacidosis (DKA) can occur if pump fails
- 24-hour telephone support may prevent acute diabetes adverse events such as DKA
- Low socioeconomic status (SES) is associated with pediatric diabetes management and outcomes
- In Ontario, low-income children are less likely to use a pump
 - Shulman, R et al. *Pediatr Diabetes*. 2016 Jan 8



+ Objectives and Hypothesis



- To explore the relationship between 24-hour support, SES and
 - 1) The risk of DKA or death
 - 2) The rate of diabetes-related admissions and emergency department (ED) visits
- We hypothesized that lack of 24-hour support and lower SES would be associated with an increased risk of adverse events




Study Design and Data Sources



- Observational population-based cohort study
- All individuals <19 years with pump funding (2006-2011) followed until 2013
- PDN centre survey (2012)
 - Centre characteristics and resources
- Administrative Data (Institute for Clinical and Evaluative Sciences (ICES))
 - Assistive devices program (ADP)
 - Ontario Diabetes Database (ODD)
 - Hospitalizations and ED visits
 - Population and Demographic Data

+ Analyses




- Cox proportional hazards model: association between time to DKA or death and 24 hour support and deprivation
- Poisson model: association between the rate of diabetes-related ED visits or admissions and 24 hour support and deprivation
- Other variables
 - Centre-level: centre type, model of physician care, nursing patient load
 - Patient-level: Prior DKA, age, duration of T1D, baseline glycemic control, comorbid condition, and rurality
- Generalized estimating equations (GEE) used to account for patient clustering at centres

+ Baseline characteristics

	24-hour support available (n=1463)	No 24-hour support available (n=1730)
Deprivation Quintile, n (%)		
1 (least deprived)	529 (36.2%)	544 (31.5%)
2	401 (27.4%)	460 (26.6%)
3	257 (17.6%)	340 (19.7%)
4	164 (11.2%)	223 (12.9%)
5 (most deprived)	102 (7.0%)	150 (8.7%)
Missing	10 (0.7%)	13 (0.8%)
At least one DKA admission in 2 years prior to index date		
Yes	139 (9.5%)	252 (14.6%)
Centre type, n (%)		
Large community	748 (51.1%)	598 (34.6%)
Small community	72 (4.9%)	570 (33.0%)
Tertiary	643 (44.0%)	562 (32.5%)
Physician model, n (%)		
Generalist	323 (22.1%)	283 (16.4%)
Pediatric Endocrinologist	965 (66.0%)	981 (56.7%)
Visiting pediatric endocrinologist	175 (12.0%)	466 (26.9%)


+ DKA or Death



- Mean time on pump was 3.77 years (SD 1.64)
- DKA rate: 5.28/100 person-years
 - Baseline DKA rate of 6.97/100 person-years
- <6 deaths
 - Death rate: 0.033/100 person-years



Cumulative Probability of DKA or death according to time since pump start



+ Time to first DKA or death (n=3193)

	DKA or death	
	Adjusted Hazard Ratio (95% CI)	p-value
Deprivation Quintile		
1 (least deprived) (n=1073)	(Ref) 1.0	
2 (n=861)	1.53 (1.22-1.91)	0.0002
3 (n=597)	0.96 (0.67-1.36)	0.8053
4 (n=387)	1.45 (1.07-1.95)	0.0161
5 (most deprived) (n=252)	1.58 (1.05-2.38)	0.0272
Missing (n=23)	1.87 (0.96-3.62)	0.0654
24 hr support		
No (n=1730)	(Ref) 1.0	
Yes (n=1463)	0.90 (0.73-1.10)	0.3023

+ Time to first DKA or death (n=3193)

	DKA or death	
	Adjusted Hazard Ratio (95% CI)	p-value
At least one DKA admission in 2 years prior to index date		
No (n=2802)	(Ref) 1.0	
Yes (n=391)	1.63 (1.28-2.07)	<0.0001
Baseline HbA1c		
<7.5% (n=718)	(Ref) 1.0	
7.5-9.0% (n=1845)	1.69 (1.31-2.18)	<0.0001
>9.0% (n=630)	2.63 (1.89-3.67)	<0.0001
Age group		
<6 th birthday (n=224)	0.88 (0.57-1.36)	0.5569
6 th -13 th birthday (n=1448)	(Ref) 1.0	
≥13 th birthday (n=1521)	2.35 (1.90-2.90)	<0.0001
Annual nursing patient load		
<200 (n=1283)	(Ref) 1.0	
≥200 (n=1910)	1.24 (1.00-1.52)	0.0468



Diabetes-related admissions or ED visits



- Rate of diabetes-related admissions and ED visits: 14.73/100 person-years
 - Baseline rate: 27.59/100 person years
- Being more deprived is associated with a higher rate of events (HR 1.60 (95% CI 1.27-2.00), $p < 0.0001$)
- 24-hour support was not significantly associated (HR 0.86 (95% CI 0.70-1.05), $p = 0.1265$)




Discussion: Adverse event rate



- Rate of DKA in Ontario pediatric pump users similar to other countries
 - Sweden (2000) (3.6/100 patient-years)
 - Hanas R. et al. *Pediatr Diabetes* 2009; 10:33-37
 - Europe and Israel (2004-2005) (6.26 /100 patient-years)
 - Danne T. et al. *Diabetologia* 2008; 51:1594-1601
- Risk DKA or death in first 2 years of pump use is low and rate of adverse events after pump start is lower compared to baseline




Discussion: 24 hour support



- Our results question previous work
 - 24-hour telephone care for T1D:
 - Prevents ED and office visits
 - Allen HF, et al. *Pediatr Diabetes* 2002; 3:95-100
 - Increases ED visits but decreases likelihood of hospitalization
 - Franklin BE, et al. *Diabetes Care* 2014; 37:81-87
- 24-hour support may not be an important service or barriers to its effective use may exist




Discussion: SES



- Our results support existing evidence for SES disparity in diabetes management and outcomes
- US Type 1 Diabetes Exchange clinic registry (n=13,487), low income is associated with a higher frequency of DKA
 - Cengiz, E. et al. *Pediatr Diabetes* 2013; 14:447-454
- In Ontario, children of lower SES are less likely to be using pumps
 - Shulman, R. et al. *Pediatr Diabetes*. 2016 Jan 8
- Single Ontario center: no association between deprivation and glycemic control in pump users
 - Zuijdwijk, C.S. et al. *J Pediatr* 2013; 162:730-735


+ Limitations



- We measured applications for pump funding, not actual pump use
- Some individuals who applied for pump funding may have been using a pump prior to universal funding
- HbA1c data is unavailable
- Self-reported nature of the centre data



Conclusions



- Implementation in Ontario not resulted in safety concerns
- Need to examine potential barriers and benefits of 24 hour support
 - Impact on parental anxiety
 - Reduction in disparity in SES of diabetes outcomes
- Need to further investigate underlying causes of SES disparities in the risk of DKA
 - Inform interventions aimed at reducing the inequity
- Targeted interventions for high risk groups may be beneficial

+ Acknowledgements

- Canadian Child Health Clinician Scientist Program (CCHCSP)
- Department of Pediatrics, the Hospital for Sick Children
- Funding from:
 - Creative Professional Activities grant from the Department of Pediatrics, Hospital for Sick Children
 - Dr. Guttman's Canadian Institute for Health Research Applied Chair in Child Health Services and Policy Research award
- Thesis committee
 - Dr. Astrid Guttman
 - Dr. Fiona Miller
 - Dr. Therese Stukel
 - Dr. Denis Daneman



+ Disclaimer

- This study was supported by the Institute for Clinical Evaluative Sciences (ICES), which is funded by an annual grant from the Ontario Ministry of Health and Long-Term Care (MOHLTC). The opinions, results and conclusions reported in this paper are those of the authors and are independent from the funding sources. No endorsement by ICES or the MOHLTC is intended or should be inferred. These datasets were linked using unique encoded identifiers and analyzed at the Institute for Clinical Evaluative Sciences (ICES).
- This study was approved by the institutional review boards at Sunnybrook Health Sciences Centre, Toronto, Canada, the University of Toronto, and The Hospital for Sick Children.
- Parts of this material are based on data and information compiled and provided by CIHI. However, the analyses, conclusions, opinions and statements expressed herein are those of the author, and not necessarily those of CIHI.

